

Anticipating PTSD in severe COVID survivors: the case for screen-and-treat

Talya Greene, Sharif El-Leithy, Jo Billings, Idit Albert, Jennifer Birch, Mari Campbell, Kim Ehntholt, Lorna Fortune, Nicola Gilbert, Nick Grey, Laurinne Hana, Helen Kennerley, Deborah Lee, Sarah Lunn, Dominic Murphy, Mary Robertson, Dorothy Wade, Chris R. Brewin & Michael A. P. Bloomfield

To cite this article: Talya Greene, Sharif El-Leithy, Jo Billings, Idit Albert, Jennifer Birch, Mari Campbell, Kim Ehntholt, Lorna Fortune, Nicola Gilbert, Nick Grey, Laurinne Hana, Helen Kennerley, Deborah Lee, Sarah Lunn, Dominic Murphy, Mary Robertson, Dorothy Wade, Chris R. Brewin & Michael A. P. Bloomfield (2022) Anticipating PTSD in severe COVID survivors: the case for screen-and-treat, *European Journal of Psychotraumatology*, 13:1, 1959707, DOI: [10.1080/20008198.2021.1959707](https://doi.org/10.1080/20008198.2021.1959707)

To link to this article: <https://doi.org/10.1080/20008198.2021.1959707>



© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 26 Jan 2022.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)



Anticipating PTSD in severe COVID survivors: the case for screen-and-treat

Talya Greene^{a,b,*}, Sharif El-Leithy^{c,*}, Jo Billings^d, Idit Albert^{d,e}, Jennifer Birch^{f,g}, Mari Campbell^h, Kim Ehntholtⁱ, Lorna Fortune^{j,k}, Nicola Gilbert^l, Nick Grey^{l,m}, Laurinne Hana^{l,n}, Helen Kennerley^{o,p}, Deborah Lee^q, Sarah Lunn^r, Dominic Murphy^e, Mary Robertsonⁱ, Dorothy Wade^s, Chris R. Brewin^t and Michael A. P. Bloomfield^{u,v}

^aDepartment of Community Mental Health, University of Haifa, Haifa, Israel; ^bDivision of Psychiatry, Institute of Mental Health, University College London, London, UK; ^cTraumatic Stress Service, South West London & St George's Mental Health NHS Trust, London, UK; ^dCentre for Anxiety Disorders and Trauma, South London & Maudsley NHS Trust, London, UK; ^eDepartment of Psychology, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK; ^fDepartment of Clinical Psychology and Psychology Therapies, Norwich Medical School, University of East Anglia, Norwich, UK; ^gNorfolk and Norwich University Hospitals NHS Foundation Trust, Norwich, UK; ^hRoyal Free London NHS Foundation Trust, London, UK; ⁱThe Traumatic Stress Clinic, St Pancras Hospital, Camden & Islington NHS Foundation Trust, London, UK; ^jBarnet Enfield & Haringey MHT, London, UK; ^kNorth Middlesex University Hospital NHS Trust, London, UK; ^lSussex Partnership NHS Foundation Trust, Sussex, UK; ^mSchool of Psychology, University of Sussex, Brighton, UK; ⁿUniversity College London Hospitals NHS Foundation Trust, London, UK; ^oOxford Cognitive Therapy Centre, Warneford Hospital, Oxford Health NHS Foundation Trust, Oxford, UK; ^pOxford Cognitive Therapy Centre, University of Oxford, Oxford, UK; ^qBerkshire Traumatic Stress Service, Berkshire Foundation NHS Trust, Reading, UK; ^rWhittington Health NHS Trust, London, UK; ^sCritical Care Department, University College London Hospitals NHS Foundation Trust, London, UK; ^tDepartment of Clinical, Educational and Health Psychology, University College London, London, UK; ^uTranslational Psychiatry Research Group, Department of Mental Health Neuroscience, Division of Psychiatry, Institute of Mental Health, University College London, London, UK; ^vNational Institute for Health Research, University College London Hospitals Biomedical Research Centre, London, UK

ABSTRACT

Based on research from previous pandemics, studies of critical care survivors, and emerging COVID-19 data, we estimate that up to 30% of survivors of severe COVID will develop PTSD. PTSD is frequently undetected across primary and secondary care settings and the psychological needs of survivors may be overshadowed by a focus on physical recovery. Delayed PTSD diagnosis is associated with poor outcomes. There is a clear case for survivors of severe COVID to be systematically screened for PTSD, and those that develop PTSD should receive timely access to evidence-based treatment for PTSD and other mental health problems by multidisciplinary teams.

Anticipando el TEPT en los sobrevivientes de COVID severo: El caso para detección y tratamiento

Basados en la investigación de pandemias previas, los estudios de los sobrevivientes a cuidados críticos, y los datos emergentes de COVID-19, estimamos que hasta un 30% de los sobrevivientes del COVID grave desarrollarán TEPT. El TEPT es frecuentemente subdetectado en los servicios de salud primarios y secundarios y las necesidades psicológicas de los sobrevivientes puede verse eclipsadas por un enfoque en la recuperación física. El diagnóstico tardío de TEPT se asocia con pobres resultados. Existe un caso claro para que los sobrevivientes del COVID grave sean evaluados sistemáticamente para detectar el TEPT, y aquellos que desarrollan un TEPT deben tener acceso oportuno a tratamientos basados en la evidencia para el TEPT y para otros problemas de salud mental por equipos multidisciplinarios.

预测重症 COVID 幸存者的 PTSD: 筛查和治疗的案例

根据先前对疫情, 重症监护幸存者的研究和新兴的 COVID-19 数据, 我们估计多达 30% 的重症 COVID 幸存者将发展出 PTSD。PTSD 经常不会被初级和二级医疗机构发现, 幸存者的心理需求可能会被对身体康复的关注所掩盖。延迟的 PTSD 诊断与不良结果相关。有一个明确的案例, 重症 COVID 幸存者需要系统性地接受 PTSD 筛查, 那些发展出 PTSD 的人应该能够及时获得多学科团队针对 PTSD 和其他心理健康问题的循证治疗。

ARTICLE HISTORY

Received 20 June 2021
Accepted 27 June 2021

KEYWORDS

Psychological trauma; mental health screening; PTSD assessment; long COVID; intensive care; critical care

PALABRAS CLAVE

Trauma Psicológico; Detección en Salud Mental; Evaluación de TEPT; COVID Largo; Cuidados Intensivos; Cuidado Crítico

关键词

心理创伤; 心理健康筛查; PTSD评估; 长期COVID; 重症监护; 危重护理

HIGHLIGHTS

- We anticipate that up to 30% of survivors of severe COVID will develop PTSD, yet PTSD is frequently undetected in primary and secondary care settings.
- There is, therefore, a clear case for establishing systematic screening and ensuring timely access to treatment.

1. Introduction

The coronavirus pandemic continues to have profound deleterious effects on public mental health (Javakhishvili et al., 2020; Pfefferbaum & North,

2020). Here we focus on how to meet the needs of survivors of severe COVID who are at especially high risk of post-traumatic stress disorder (PTSD). Severe COVID illness is life-threatening and its clinical management can be highly invasive and frightening

CONTACT Michael A. P. Bloomfield  m.bloomfield@ucl.ac.uk  Translational Psychiatry Research Group, Department of Mental Health Neuroscience, Division of Psychiatry, Institute of Mental Health, University College London, Maple House, 149 Tottenham Court Road, London W1T 7NF, UK

*Denotes equal contribution as first author.

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

(Rogers et al., 2020). Surviving severe COVID therefore fulfils internationally-recognized definitions of exposure to psychological trauma and is thus a risk factor for PTSD (Bisson, Cosgrove, Lewis, & Roberts, 2015; World Health Organization, 2018). Here we describe an emerging strategy of 'screen-and-treat' programmes for PTSD after severe COVID – which involve coordinated proactive outreach, assessment, and treatment. These programmes have been implemented after various kinds of major incidents such as terrorist attacks, and are designed to systematically screen and provide evidence-based treatment to those who need it. We argue that there is a critical and immediate need to integrate PTSD screen-and-treat into the follow-up care of patients recovering from severe COVID, including those survivors with long COVID.

2. PTSD

PTSD is a potentially debilitating disorder that can develop after exposure to exceptionally threatening events. PTSD is associated with high rates of comorbidity of mental and physical illness (APA, 2013). The core symptoms of PTSD include intrusive memories of the trauma, avoidance of trauma-related stimuli, negative alterations in cognitions and mood, and a persistent sense of threat and hyperarousal (Bisson et al., 2015; World Health Organization, 2018). PTSD is a treatable condition using evidence-based interventions including psychotherapies such as trauma-focused cognitive therapy and eye movement desensitization and reprocessing (EMDR) (APA, 2013). There is also an evidence base for pharmacological treatments (Cipriani et al., 2018).

Yet in routine clinical practice, PTSD is underdiagnosed (Greene, Neria, & Gross, 2016), even in specialist psychiatric settings (Zammit et al., 2018). Delayed diagnosis of PTSD is associated with poorer clinical outcomes and comorbidity (Priebe et al., 2009). It is therefore essential that services are planned in such a way to overcome these barriers to diagnosis.

3. Severe COVID is a risk factor for PTSD

Several lines of evidence indicate that severe COVID illness is a risk factor for PTSD. Research with survivors of past coronavirus disease outbreaks, including Severe Acute Respiratory Syndrome (SARS) and Middle East Respiratory Syndrome (MERS), found that PTSD rates were high following hospital discharge, and remained elevated for months or even years (Salehi et al., 2021). A recent study of 381 survivors of severe COVID found that 30.2% were diagnosed with PTSD (Janiri et al., 2021). Another study found that hospitalized COVID patients were likewise at increased risk of PTSD (Chamberlain, Grant, Trender, Hellyer, & Hampshire, 2021). PTSD

symptoms were highest among those who had been ventilated; 35% of those who had been ventilated endorsed all ten of the PTSD symptoms measured in the study. We also know from research prior to COVID that around 20%-25% of people discharged from critical care report symptoms of PTSD (Parker et al., 2015).

There are particular aspects of the experience of critically ill COVID patients that are likely to increase the risk of traumatization and therefore PTSD (Kaseda & Levine, 2020). These biopsychosocial risk factors are related directly to the impact of the illness, as well as to aspects of its treatment, and the context within which these occur. High rates of delirium have been reported among COVID patients due to infection, sedation, and the clinical environment (Kaseda & Levine, 2020). Delirium is often associated with terrifying experiences including hallucinations and delusions, and these experiences can be psychologically traumatic in themselves (Wade et al., 2015). Furthermore, the breathing difficulties commonly experienced by patients with COVID are particularly horrific and have been described by patients as the prolonged sensation of drowning (Ferguson et al., 2020). In addition, COVID patients are sedated more deeply and ventilated for longer than typical for critical care treatment (Ferguson et al., 2020), which is associated with a further elevation in risk of PTSD beyond that already observed in critical care patients.

Multiple additional factors are likely to exacerbate psychological distress among COVID survivors during their acute illness, including witnessing the deaths of other patients, isolation from loved ones, and the use of personal protective equipment by staff which may interfere with normal communication and can feel impersonal. In addition, following discharge from hospital, the normal recovery process from a psychologically traumatic event may be impaired by a range of factors associated with the pandemic. COVID survivors may experience continued isolation from loved ones and their other usual sources of support due to community infection control measures and ongoing social restrictions. Survivors may have been bereaved by COVID or have family members who are also seriously ill. Many survivors recover slowly and continue to suffer longer-term physical effects of their illness for many months. All of this takes place within the wider social, political and economic context of uncertainty, restrictions and physical distancing, which will exacerbate psychological distress, further contributing to a sense of ongoing threat and therefore impaired recovery.

Taken together, there is a coherent body of evidence that severe COVID increases the risk of PTSD. Based on literature from previous pandemics, studies of critical care survivors, and emerging COVID-19 studies, we estimate that up to 30% of survivors of severe COVID-19 will develop PTSD. These rates are similar to survivors of mass casualty events such as transport disasters and terrorist attacks. While definitive epidemiological

evidence will take time to arrive, we cannot wait for this to begin planning for the assessment and treatment of these patients as there is already urgent clinical need right now.

4. Challenges in detecting PTSD

The COVID pandemic is a mass casualty incident. Previous experience with mass casualty incidents internationally tells us that PTSD in survivors is frequently not detected through conventional routes such as primary care or hospital outpatient clinics (Allsopp et al., 2019; Pfefferbaum, North, Flynn, Norris, & DeMartino, 2002). This is likely the result of multiple causes. First, mental health was not a part of most countries' official pandemic planning (Brewin, DePierro, Pirard, Vasquez, & Williams, 2020). Second, psychological needs can be overshadowed by a focus on physical recovery. In many cases primary care and general hospital healthcare staff lack the time and/or experience necessary to detect PTSD. Third, individuals often do not know how to recognize and seek help for psychological difficulties. Fourth, there is wide local variation in the provision of hospital-based critical care follow-up clinics, and in the survivor's ability to access support even when it does exist. Together, this highlights the need for urgent detection of PTSD beyond routine care through new, dedicated clinical pathways.

5. Assessing and monitoring symptoms of PTSD

The deployment of COVID-PTSD screen-and-treat programmes offers a solution to these challenges. The purposes of a COVID-PTSD screen-and-treat programme are to proactively identify, follow-up, and treat survivors following hospital admission in tandem with multidisciplinary clinics. We previously provided guidance on screen-and-treat programmes for PTSD for survivors of severe COVID illness (El-Leithy et al., 2020). Our guidance draws upon a body of research evidence and clinical guidelines derived from previous epidemics, mass casualty events, and critical care settings.

There are strategies to reduce the risk of PTSD in critical care settings (Tingey, Bentley, & Hosey, 2020), but here we focus on the detection and follow up of these patients after the acute phase of the COVID illness. We propose that brief psychological tools (i.e. validated questionnaires) are administered to COVID survivors prior to discharge from acute hospitals and again four weeks after discharge to identify possible emerging mental health needs (see Figure 1). The brief assessment at four weeks must include a screening instrument for PTSD, such as the trauma screening questionnaire (Brewin et al., 2002). Comprehensive assessments for PTSD should then be carried out when people score above clinical thresholds on those screening tools. Advice and repeated follow-up for 12 months following

discharge should be offered to people scoring below threshold to actively monitor the trajectory of their recovery and to ensure that cases of delayed onset PTSD are not missed.

6. Overcoming barriers to accessing treatment

Critically, following appropriate screening and identification of PTSD, there is a clinical and ethical imperative to provide access to evidence-based treatment in a timely manner. Yet in many healthcare systems, even in routine times, PTSD treatment provision is inadequate at best and woeful at worst. Therefore, funding should immediately be made available across health systems to urgently enable the assessment and increased capacity for treatment of PTSD in survivors of severe COVID. It is essential that these programmes are coordinated with existing specialist psychological trauma services in secondary and tertiary care that have the appropriate expertise along the spectrum of clinical severity and complexity.

The international experience from other mass casualty events shows that PTSD screen-and treat programmes are associated with improved clinical outcomes (Brewin et al., 2010; French et al., 2019; Gobin et al., 2018; Gouweloos-Trines et al., 2019). Given the potential complexity and comorbidities associated with severe COVID illness, these teams should manage PTSD as part of an integrated framework. These programmes can run alongside hospital rehabilitation and are best delivered by dedicated multidisciplinary teams with PTSD expertise, including with input from specialist psychological trauma services where available.

Severe COVID illness disproportionately affects black, ethnic minority, elderly and other disadvantaged groups. Therefore, any effective mental health response will need to be proactive in identifying, engaging with, and adapting to the specific needs of those communities most affected by COVID, in order to ensure equality of access to assessment and treatment. This adds further weight to the argument for COVID-specific PTSD assessment and treatment programmes, which will have the capability to systematically monitor outcomes and inform future responses.

7. Conclusion

There is no health without mental health. Developing local, regional, and national programmes for severe COVID survivors places their mental health needs on an equal footing to those from other mass casualty events, such as terrorist incidents. We anticipate that up to 30% of survivors of severe COVID illness will develop PTSD. Given the high prevalence of COVID across the globe, it likely that millions of people will develop PTSD as a direct result of the pandemic. PTSD is often undetected and




Figure 1. Screen-and-treat process.

untreated resulting in suffering and premature death for patients. This is especially the case when healthcare systems are over-stretched and when the focus is on the recovery from a 'physical' rather than 'mental' illness. Therefore, it is critical to anticipate and meet this projected need. Screen-and-treat programmes are known to be feasible, acceptable and currently have the most support. There is thus a clear case for survivors of severe COVID to be systematically screened for PTSD, and those that develop PTSD should receive timely access to evidence-based treatment.

Disclosure statement

No potential conflict of interest was reported by the author(s).

ORCID

Talya Greene  <http://orcid.org/0000-0002-3044-2841>
 Jo Billings  <http://orcid.org/0000-0003-1238-2440>
 Idit Albert  <http://orcid.org/0000-0002-3800-1530>
 Dominic Murphy  <http://orcid.org/0000-0002-9530-2743>
 Dorothy Wade  <http://orcid.org/0000-0001-6431-3776>
 Chris R. Brewin  <http://orcid.org/0000-0002-7462-4460>
 Michael A. P. Bloomfield  <http://orcid.org/0000-0002-1972-4610>

References

- Allsopp, K., Brewin, C. R., Barrett, A., Williams, R., Hind, D., Chitsabesan, P., & French, P. (2019). Responding to mental health needs after terror attacks. *BMJ*, 366, l4828. doi:10.1136/bmj.l4828
- APA. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: Author.
- Bisson, J. I., Cosgrove, S., Lewis, C., & Roberts, N. P. (2015). Post-traumatic stress disorder. *BMJ*, 351, h6161. doi:10.1136/bmj.h6161
- Brewin, C. R., DePierro, J., Pirard, P., Vasquez, C., & Williams, R. (2020). Why we need to integrate mental health into pandemic planning. *Perspectives in Public Health*, 140, 309–310. doi:10.1177/1757913920957365
- Brewin, C. R., Fuchkan, N., Huntley, Z., Robertson, M., Thompson, M., Scragg, P., ... Ehlers, A. (2010). Outreach and screening following the 2005 London bombings: Usage and outcomes. *Psychological Medicine*, 40, 2049–2057. doi:10.1017/S0033291710000206
- Brewin, C. R., Rose, S., Andrews, B., Green, J., Tata, P., McEvedy, C., ... Foa, E. B. (2002). A brief screening instrument for posttraumatic stress disorder. *British Journal of Psychiatry*, 181, 158–162. doi:10.1192/bjp.181.2.158
- Chamberlain, S., Grant, J. E., Trender, W., Hellyer, P., & Hampshire, A. (2021). Post-traumatic stress disorder (PTSD) symptoms in COVID-19 survivors: Online population survey. *British Journal of Psychiatry Open*, 7. doi:10.1192/bjo.2021.3
- Cipriani, A., Williams, T., Nikolakopoulou, A., Salanti, G., Chaimani, A., Ipser, J., ... Stein, D. J. (2018). Comparative efficacy and acceptability of pharmacological treatments for post-traumatic stress disorder in adults: A network meta-analysis. *Psychological Medicine*, 48, 1975–1984. doi:10.1017/S003329171700349X
- El-Leithy, S., Albert, I., Billings, J., Birch, J., Robertson, M., Brewin, C. R., ... Bloomfield, M. (2020, October 17). Guidance on screening and active monitoring for post-traumatic stress disorder (PTSD) and other mental health consequences in people recovering from severe COVID-19 illness. COVID Trauma Response Working Group. Retrieved from https://232fe0d6-f8f4-43eb-bc5d-6aa50ee47dc5.filesusr.com/ugd/6b474f_733283bf71ce4295b915dffa86886280.pdf
- Ferguson, N., Laydon, D., Nedjati Gilani, G., Imai, N., Ainslie, K., Baguelin, M., ... Dighe, A. (2020). *Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID19 mortality and healthcare demand*. London, UK: WHO Collaborating Centre for Infectious Disease Modelling, MRC Centre for Global Infectious Disease Analysis, Abdul Latif Jameel Institute for Disease and Emergency Analytics, and Imperial College London. doi:10.25561/77482.
- French, P., Barrett, A., Chitsabesan, P., Stancombe, J., Allsopp, L. K., Brewin, C. R., ... Deol, J. (2019). Psychological screening of adults and young people following the Manchester Arena incident. *British Journal of Psychiatry Open*, 5, e85. doi:10.1192/bjo.2019.61
- Gobin, M., Rubin, G. J., Albert, I., Beck, A., Danese, A., Greenberg, N., ... Oliver, I. (2018). Outcomes of mental health screening for UK Nationals affected by the 2015–

- 2016 terrorist attacks in Tunisia, Paris, and Brussels. *Journal of Traumatic Stress*, 31, 471–479. doi:10.1002/jts.22317
- Gouweloos-Trines, J., te Brake, H., Sijbrandij, M., Boelen, P. A., Brewin, C. R., & Kleber, R. J. (2019). Evaluating a psychosocial support program for survivors of an airplane crash: Screening for PTSD and depression and assessment of self-reported treatment needs. *European Journal of Psychotraumatology*, 10. doi:10.1080/20008198.2018.1554406
- Greene, T., Neria, Y., & Gross, R. (2016). Prevalence, detection and correlates of PTSD in the primary care setting: A systematic review. *Journal of Clinical Psychology in Medical Settings*, 23, 160–180. doi:10.1007/s10880-016-9449-8
- Janiri, D., Carfi, A., Kotzalidis, G. D., Bernabei, R., Landi, F., & Sani, G. (2021, February 18). Gemelli against COVID-19 Post-Acute Care Study Group. Posttraumatic stress disorder in patients after severe COVID-19 infection. *JAMA Psychiatry*, 78, 567. doi:10.1001/jamapsychiatry.2021.0109
- Javakhishvili, J. D., Ardino, V., Bragesjö, M., Kazlauskas, E., Olf, M., & Schäfer, I. (2020). Trauma-informed responses in addressing public mental health consequences of the COVID-19 pandemic: Position paper of the European Society for Traumatic Stress Studies (ESTSS). *European Journal of Psychotraumatology*, 11 (1), 1780782. doi:10.1080/20008198.2020.1780782
- Kaseda, E. T., & Levine, A. J. (2020). Post-traumatic stress disorder: A differential diagnostic consideration for COVID-19 survivors. *The Clinical Neuropsychologist*, 34, 1498–1514. doi:10.1080/13854046.2020.1811894
- Parker, A. M., Sricharoenchai, T., Raparla, S., Schneck, K. W., Bienvenu, O. J., & Needham, D. M. (2015). Posttraumatic stress disorder in critical illness survivors: A meta-analysis. *Critical Care Medicine*, 43, 1121–1129. doi:10.1097/CCM.0000000000000882
- Pfefferbaum, B., & North, C. S. (2020). Mental health and the Covid-19 pandemic. *New England Journal of Medicine*, 383, 510–512. doi:10.1056/NEJMp2008017
- Pfefferbaum, B., North, C. S., Flynn, B. W., Norris, F. H., & DeMartino, R. (2002). Disaster mental health services following the 1995 Oklahoma City bombing: Modifying approaches to address terrorism. *CNS Spectrums*, 7, 575–579. doi:10.1017/S1092852900018174
- Priebe, S., Matanov, A., Janković Gavrilović, J., McCrone, P., Ljubotina, D., Knežević, G., ... Schützwoh, M. (2009). Consequences of untreated post-traumatic stress disorder following war in former Yugoslavia: Morbidity, subjective quality of life, and care costs. *Croatian Medical Journal*, 50, 465–475. doi:10.3325/cmj.2009.50.465
- Rogers, J. P., Chesney, E., Oliver, D., Pollak, T. A., McGuire, P., Fusar-Poli, P., & David, A. S. (2020). Psychiatric and neuropsychiatric presentations associated with severe coronavirus infections: A systematic review and meta-analysis with comparison to the COVID-19 pandemic. *The Lancet Psychiatry*, 7, 611–627. doi:10.1016/S2215-0366(20)30203-0
- Salehi, M., Amanat, M., Mohammadi, M., Salmanian, M., Rezaei, N., Saghazadeh, A., & Garakani, A. (2021). The prevalence of post-traumatic stress disorder related symptoms in coronavirus outbreaks: A systematic-review and meta-analysis. *Journal of Affective Disorders*, 282, 527–538. doi:10.1016/j.jad.2020.12.188
- Tingey, J. L., Bentley, J. A., & Hosey, M. M. (2020). COVID-19: Understanding and mitigating trauma in ICU survivors. *Psychological Trauma: Theory, Research, Practice, and Policy*, 12, S100. doi:10.1037/tra0000884
- Wade, D. M., Brewin, C. R., Howell, D. C., White, E., Mythen, M. G., & Weinman, J. A. (2015). Intrusive memories of hallucinations and delusions in traumatized intensive care patients: An interview study. *British Journal of Health Psychology*, 20, 613–631. doi:10.1111/bjhp.12109
- World Health Organization. (2018). *International Statistical classification of diseases and related health problems* (11th Rev.). Geneva, Switzerland: Author.
- Zammit, S., Lewis, C., Dawson, S., Colley, H., McCann, H., Piekarski, A., ... Bisson, J. (2018). Undetected post-traumatic stress disorder in secondary-care mental health services: Systematic review. *The British Journal of Psychiatry*, 212, 11–18. doi:10.1192/bjp.2017.8